

## Maternal effector T cells within decidua:

Lissauer, D.; Kilby, M. D.; Moss, P.

DOI:

[10.1016/j.placenta.2017.09.003](https://doi.org/10.1016/j.placenta.2017.09.003)

License:

Creative Commons: Attribution-NonCommercial-NoDerivs (CC BY-NC-ND)

*Document Version*

Peer reviewed version

*Citation for published version (Harvard):*

Lissauer, D, Kilby, MD & Moss, P 2017, 'Maternal effector T cells within decidua: The adaptive immune response to pregnancy?', *Placenta*, vol. 60, pp. 140-144. <https://doi.org/10.1016/j.placenta.2017.09.003>

[Link to publication on Research at Birmingham portal](#)

### General rights

Unless a licence is specified above, all rights (including copyright and moral rights) in this document are retained by the authors and/or the copyright holders. The express permission of the copyright holder must be obtained for any use of this material other than for purposes permitted by law.

- Users may freely distribute the URL that is used to identify this publication.
- Users may download and/or print one copy of the publication from the University of Birmingham research portal for the purpose of private study or non-commercial research.
- User may use extracts from the document in line with the concept of 'fair dealing' under the Copyright, Designs and Patents Act 1988 (?)
- Users may not further distribute the material nor use it for the purposes of commercial gain.

Where a licence is displayed above, please note the terms and conditions of the licence govern your use of this document.

When citing, please reference the published version.

### Take down policy

While the University of Birmingham exercises care and attention in making items available there are rare occasions when an item has been uploaded in error or has been deemed to be commercially or otherwise sensitive.

If you believe that this is the case for this document, please contact [UBIRA@lists.bham.ac.uk](mailto:UBIRA@lists.bham.ac.uk) providing details and we will remove access to the work immediately and investigate.

# Accepted Manuscript

Maternal effector T cells within decidua: The adaptive immune response to pregnancy?

D. Lissauer, M.D. Kilby, P. Moss



PII: S0143-4004(17)31107-4

DOI: [10.1016/j.placenta.2017.09.003](https://doi.org/10.1016/j.placenta.2017.09.003)

Reference: YPLAC 3724

To appear in: *Placenta*

Received Date: 1 May 2017

Revised Date: 25 August 2017

Accepted Date: 7 September 2017

Please cite this article as: Lissauer D, Kilby MD, Moss P, Maternal effector T cells within decidua: The adaptive immune response to pregnancy?, *Placenta* (2017), doi: 10.1016/j.placenta.2017.09.003.

This is a PDF file of an unedited manuscript that has been accepted for publication. As a service to our customers we are providing this early version of the manuscript. The manuscript will undergo copyediting, typesetting, and review of the resulting proof before it is published in its final form. Please note that during the production process errors may be discovered which could affect the content, and all legal disclaimers that apply to the journal pertain.

# **Maternal effector T cells within decidua: the adaptive immune response to pregnancy?**

**Lissauer D<sup>1</sup>, Kilby, MD<sup>1</sup>. and Moss, P<sup>2, 3</sup>**

<sup>1</sup> Centre for Women's and Newborn's Health and Centre of Endocrinology, Diabetes and Metabolism [CEDAM], College of Medical and Dental Sciences, University of Birmingham and Birmingham Health Partners, UK

<sup>2</sup> Institute of Immunology and Immunotherapy, College of Medical and Dental Sciences, University of Birmingham and Birmingham Health Partners, UK

<sup>3</sup> Corresponding Author. p.moss@bham.ac.uk

**Abstract**

In human pregnancy the maternal immune system plays a critical role in the regulation of many aspects of human reproduction including implantation, placentation and defence against infection. Interest has been focussed on the role of uterine natural killer cells (uNK) in the maternal decidua whereas effector CD4+ and CD8+ T cells have received much less attention despite the observation that they represent a major proportion of decidual leucocytes in the latter phase of pregnancy. A range of recent studies have demonstrated that human decidual T cells are highly differentiated, express a range of cytokines and cytotoxic markers, and demonstrate a unique transcriptional profile characterised by high level expression of genes involved in interferon-signalling. Moreover, subpopulations of effector T cells demonstrate specificity for fetal tissue and are regulated through expression of inhibitory checkpoint proteins and T regulatory cells. Nevertheless, many questions remain to be answered, such as the potential role of maternal effector T cells in either supporting successful pregnancy or potentially clearing fetal cells that have entered the maternal circulation. In addition, there is an increasing interest in the role of maternal effector T cells in the pathogenesis of disorders such as chronic villitis, miscarriage, stillbirth, fetal growth restriction and pre-eclampsia. Current debates in relation to these questions will be discussed within this review.

## **Maternal effector T cells comprise the major population of leucocytes within decidua by the end of pregnancy**

The maintenance of a semi-allogeneic fetus within the mother represents a considerable challenge to the maternal immune system during pregnancy [1]. A wide range of mechanisms have been postulated as being important and have evolved in order to limit maternal immune recognition of fetal tissue. One of the most straightforward approaches might have been to exclude effector CD4+ and CD8+ T cells from the decidual bed. However, effector T cells represent around 60% of the T cell pool in the later stages of pregnancy [2] and whilst NK cells are relatively more numerous in the early stages of pregnancy their numbers remain stable whilst those of T cells show a gradual increase with advancing gestation [2,3]. As such this temporal replacement of NK cells by T cells mirrors the kinetics of a peripheral adaptive immune response and the uterine environment may perhaps be seen to recapitulate short term lymphocyte dynamics over a 9 month period [4]. CD45RO+ effector T cells comprise around 60% of the decidual T cell repertoire at term whilst representing only 30% of T cells within blood. This relative increase in 'antigen-experienced' T cells could potentially indicate evidence of local activation although it may also reflect selective recruitment of effector cells into decidual tissue. In this regard it is important to compare the phenotypic and functional features of T cells in the two compartments and this is shown in Table 1. This reveals a number of differences between effector cells in decidua and blood, including the observation that decidual effector cells are more highly differentiated than peripheral cells, with over 40% of such cells demonstrating a CD27-CD28- phenotype compared to less than 20% in blood [5].

This pattern of local activation of decidual T cells might be taken to represent recognition of fetal tissue. Fetal trophoblast cells downregulate the expression of HLA-A and HLA-B, and observation that mismatch of polymorphic HLA-C alleles between mother and fetus was associated with increased levels of T cell activation provided some of the first support for this hypothesis [6].

## **The CXCL10-CXCR3 axis is important in attracting effector T cells to decidual tissue**

Chemokines are important regulators of leukocyte migration and are therefore likely to play a major role in reproductive biology. CXCR3 is a receptor for inflammatory chemokines and studies within pregnant mice have shown that it undergoes epigenetic silencing on T cells within decidua [7], although this may be overcome in the setting of local inflammation [8]. In contrast our findings reveal CXCR3 to be expressed on 17% of human decidual CD4<sup>+</sup> T cells, one of many differences found between the immunological environment between mice and humans [5]. CXCL10, an important ligand for CXCR3, is strongly expressed and identifies the CXCL10-CXCR3 axis as an important mediator of effector cell migration into decidual tissue.

### **Maternal effector T cells within decidua can recognize fetal tissue**

It is now clear that an adaptive immune response against fetal tissue develops in most, if not all, human pregnancies. This is indicated most clearly in the development of alloreactive HLA-specific antibodies and the development of sensitive assays has revealed that these are found in most mothers following a single pregnancy [9]. Moreover, the presence of IgG isotypes reveals that antigen-specific T cell 'help' has also been established. The potential importance of these antibodies in relation to fetal health is uncertain and such responses may simply reflect an epiphenomenon that is of no consequence to pregnancy outcome. However some studies do reveal a weak clinical association and HLA-C specific antibodies have been shown to be more common in women with recurrent fetal loss (miscarriage) [10].

Given this observation, it is perhaps not surprising that maternal cellular adaptive immune responses against fetal tissue are also generated during human pregnancy. Culture and expansion of maternal T cells *in vitro* first identified cells that were able to recognise paternal cells [11] and these observations have been substantiated by techniques that directly visualise T cells with alloreactive potential. Indeed our own work [12, 13], and that of others [14-16], has identified HY-specific CD8<sup>+</sup> T cells in the maternal circulation following male pregnancies. These CD8 T cells are present for many years following pregnancy and can be reactivated *in vitro* to generate highly cytotoxic T cells that lyse male cells. HY-specific CD8<sup>+</sup> cells were detectable in 32% of women following a single male pregnancy and this proportion rose to 50% of those with 2 or more male pregnancies, indicating that alloreactive cellular immunity is boosted by recurrent episodes of fetal microchimerism. Until recently, it has not been possible to directly identify T cells with fetal specificity within decidual tissue but

we have also utilized HLA-peptide multimer technology to identify HY-specific CD8 T cells in decidual tissue [5]. Indeed, the frequency of such cells is greatly increased compared to peripheral blood and indicates that cytotoxic cells with specificity for fetal tissue are localised in direct anatomical contact.

### **Effector T cells within decidua display a novel profile of functional activity**

Tilbergs et al studied CD8<sup>+</sup> effector T cells within decidual tissue and demonstrated a unique Th1 pattern of high level IFN $\gamma$  expression together with low levels of perforin and granzyme [17]. Our own studies of decidual CD4 and CD8<sup>+</sup> T cells confirmed IFN $\gamma$  expression in many cells but also revealed expression of IL-4 in a minority population. In particular whilst IFN $\gamma$  expression was observed in 60% of CD8<sup>+</sup> T cells, 1.2% of cells also expressed IL-4, a value which, whilst relatively modest, was higher than expression within 0.7% of CD8<sup>+</sup> cells within maternal peripheral blood. Comparable values for CD4<sup>+</sup> T cells were 25% and 5% respectively and IL-4 expression was markedly higher than on maternal T cells from peripheral blood [5]. Interestingly, IL-4 expression can be induced from peripheral T cells following incubation with progesterone and this 'T<sub>prog</sub>' phenotype may therefore partially reflect the effect of the local hormonal microenvironment [18, 19].

We recently completed a comparative transcriptional analysis of effector CD4 and CD8<sup>+</sup> T cells from decidua and maternal peripheral blood. A wide range of genes were differentially expressed in decidual T cells with a striking upregulation of those which encode proteins involved in the signalling response to interferon [5]. This profile is highly unusual within effector T cells and suggests that decidual tissue is characterized by high levels of local interferon production. This subject has been relatively poorly studied although immunohistochemical expression of type 1 interferon has been observed within cells of the monocytic lineage [20,21]. It is interesting to speculate on what may drive interferon production but it is well established that endogenous retroviruses play an important role in the generation of syncytiotrophoblast [22-26]. Notably, the *Syncytin-1* protein encoded from the *env* gene of *ERVW-1* has an essential role in formation of the syncytiotrophoblast and is released into the periphery via placental microvesicles which are themselves able to illicit a T cell response [27].

### **Effector T cells within decidua demonstrate specificity for pathogens**

Pregnancy is associated with altered regulation of immune responses that can potentially increase susceptibility to some infectious diseases. The increased mortality rate of pregnant women following avian influenza infection is one such example. As such, it is not surprising that T cells within decidual tissue exhibit specificity for local pathogens [28]. T cells with specificity for cytomegalovirus and Epstein-Barr virus are preferentially recruited into decidua and mediate pathogen surveillance of maternal cells [29]. Importantly, these populations recognise peptides restricted by HLA-A and HLA-B alleles which are themselves not presented on fetal trophoblast. Recent investigations have also shown that effector cells can recognise peptides restricted through HLA-C alleles, potentially indicating efficacy in control of infected fetal tissue [30].

#### **The function of effector T cells is modulated through intrinsic and extrinsic regulation**

The finding of large numbers of functional effector T cells within decidual tissue raises the question of how such cells are regulated in order to limit potential immunopathology or fetal damage. In this regard extrinsic regulation mediated through T regulatory cells and cell-intrinsic expression of inhibitory checkpoint proteins are both emerging as important control mechanisms.

T regulatory cells are increased within decidual tissue at term pregnancy and can comprise over 20% of all CD4<sup>+</sup> T cells [31-33]. Indeed, murine experiments have shown that depletion of T regulatory cells can trigger fetal rejection [34] and it has been suggested that the evolution of the FoxP3<sup>+</sup> T regulatory cell was a key event in the development of eutherian reproduction [35]. In line with previous reports [33], we also find that decidual T cells proliferate in response to cord blood lymphocytes and that this is increased following depletion of T regulatory cells [5]. These observations indicate that such decidual populations do have specificity for maternal antigens and that this is at least partly regulated through the action of autologous T regulatory populations.

Checkpoint proteins such as PD-1 and TIM-3 are now considered to be amongst the most important molecules within clinical medicine [36, 37]. This is due to the dramatic efficacy of antibody-mediated blockade of PD-1 function in the treatment of solid tumours [38]. PD-1 expression on T cells is often taken to represent an



‘exhausted’ state and is believed to reflect cells that have undergone repeated stimulation within antigen. However, despite the considerable success of checkpoint blockade in cancer therapy there is less understanding as to the physiological role of checkpoint proteins in human T cell physiology. Interestingly, a high level of checkpoint protein expression is observed on T cells within decidua [39]. Wang et al reported large populations of Tim-3+PD-1+ CD8+ T cells within decidua during early human pregnancy and showed that incubation of CD8 T cells with trophoblast led to further checkpoint upregulation [40,41]. Interestingly, PD-L1 is expressed on syncytiotrophoblast, as well as intermediate trophoblastic cells located in the chorion laeve and implantation site [42], and some studies have indicated that PD-1 blockade in pregnant mice may result in fetal loss [43]. Of note, these observations provide further evidence that the study of reproductive immunology will be of huge importance in understanding tumour immunology and lend support to the concept of cancer as a ‘somatic pregnancy’ [44].

#### **Fetal-specific maternal T cells may play a role in limiting fetal microchimerism**

It is now generally accepted that pregnancy leads to a state of microchimerism within the mother in which significant amounts of fetal tissue and cells are released into the maternal circulation [45]. Moreover, these fetal cells have been shown to survive for long periods within the mother and have even been implicated in a range of clinical disorders such as thyroiditis. Whilst such fetal cells may provide potential benefit to the mother, such as potentially supporting repair of maternal tissue, it would seem reasonable that such chimerism would need to be controlled by the maternal immune system. In this regard, the humoral and cellular maternal response against fetal tissue may have an important role in the suppression of fetal chimerism. Indeed, some support for this hypothesis comes from the observation that the degree of chimerism has been reported to fall with repeated pregnancies, whilst the magnitude of fetal-specific immunity appears to increase [46]. These observations could provide important insights into the regulation of chimerism in disorders such as transplantation.

#### **Maternal T cell responses against fetal tissue may be associated with obstetric complications**

The observation that maternal T cells demonstrate antigenic specificity for fetal tissue raises the obvious question as to whether these may be implicated in the pathogenesis of pregnancy complications. Relatively little evidence exists to implicate maternal T cell responses in the development of pre-eclampsia and this may reflect the fact that the cardinal feature of impaired trophoblast invasion of maternal spiral arteries is determined early within pregnancy and at a time when natural killer cells dominate the cellular infiltrate. More recently, Leavey et al have suggested that T cell mediated pathology may indeed be related to a subset of mothers with preeclampsia who demonstrate a form of disorder characterized by poor fetal outcome and growth restriction but relatively less impact on maternal health [47].

Despite this, fetal-specific immunity is strongly implicated in the development of chronic villitis ('villitis of uncertain/unknown aetiology'; VUE) which is a relatively common cause of fetal growth restriction and pre-term delivery [48-51]. VUE is characterized by an inflammatory cell infiltrate of placental macrophages and T cells within the villi which develops in the absence of infection. A specific feature is non-uniform involvement of villi and ultimately this can lead to "obliterative fetal vasculopathy". Interestingly, this maternal infiltrate typically shows a CD8:CD4 ratio around 3:1 and T cells comprise almost half of the cellular infiltrate. It has been suggested that such villitis may represent relatively uncontrolled maternal immune recognition of fetal tissue and this is supported by the finding that antibodies against fetal tissue are commonly observed in this disorder and associate with the deposition of C4 complement components. However caution is needed in this interpretation as some degree of VUE can be observed in around 10% of pregnancies and as such this process might even be regarded as a 'normal variant'.

### **Maternal effector T cells represent a fascinating topic for future investigation**

Within a relatively short period of time conventional wisdom has moved from the belief that the placenta represents an effective barrier between the mother and fetus to an understanding that cells traffic between circulations and immune cells play an important role in supporting pregnancy. Decidual NK cells have been shown to possess a range of unique functional properties that represent novel immunotherapeutic opportunities [52] (Figure 1). Investigation into the physiological and potential immunopathological role of maternal alloreactive T cells has followed

more slowly. Despite this, it is now clear that maternal effector CD45RO<sup>+</sup> T cells represent the majority of T cells within decidual tissue in the latter stages of pregnancy and display a range of novel properties. It remains possible that the primary role of these cells lies in the control of local infection with no major significance in relation to reproductive outcome. Nevertheless, further investigation into the specificity and unique properties of these remarkable cells may uncover novel insights into the physiology of human placentation, the pathogenesis of reproductive disorders and offer clues towards a range of additional disorders such as cancer and transplantation biology.

248

249

250 **Acknowledgement**

251 Presented at the 2016 Global Pregnancy Collaboration (CoLab) conference in  
252 Oxford, England. The Global Pregnancy Collaboration is part of the Pre-eclampsia-  
253 Eclampsia Monitoring, Prevention & Treatment (PRE-EMPT) initiative funded by the  
254 University of British Columbia, a grantee of the Bill & Melinda Gates Foundation.

255

256

## References

1. Moffett, A., and C. Loke, Immunology of placentation in eutherian mammals, *Nat. Rev. Immunol.* 6: (2006) 584–594.
2. Stallmach T, Hebisch G, Orban P and Lu X.  
Aberrant positioning of trophoblast and lymphocytes in the foeto-maternal interface with pre-eclampsia. *Virchows Arch.* (1999) 434(3):207-11.
3. Sykes, L., D. A. MacIntyre, X. J. Yap, T. G. Teoh, P. R. Bennett, L. Sykes, D. A. MacIntyre, X. J. Yap, T. G. Teoh, and P. R. Bennett, The Th1:Th2 Dichotomy of Pregnancy and Preterm Labour. *Mediators Inflamm.* (2012): 1–12.
4. Williams, P. J., R. F. Searle, S. C. Robson, B. A. Innes, and J. N. Bulmer. Decidual leucocyte populations in early to late gestation normal human pregnancy. *J. Reprod. Immunol.* (2009) 82: 24–31.
5. Powell R, Lissauer D, Tamblyn J, Beggs A, Cox P, Moss P and Kilby MD. Decidual T cells exhibit a highly differentiated phenotype and demonstrate potential foetal-specificity and a strong transcriptional response to interferon  $\gamma$ . *J. Immunol.* In Press.
6. Tilburgs, T., S. A. Scherjon, B. J. van der Mast, G. W. Haasnoot, M. Versteeg-V D Voort-Maarschalk, D. L. Roelen, J. J. van Rood, and F. H. J. Claas. Foetal-maternal HLA-C mismatch is associated with decidual T cell activation and induction of functional T regulatory cells. *J. Reprod. Immunol.* (2009) 82: 148–57.
7. Nancy, P., and A. Erlebacher, Epigenetic repression of chemokine expression at the maternal-foetal interface as a mechanism of foeto-maternal tolerance. *Médecine Sci. M/S* 28: (2012) 1037–9.
8. Chaturvedi V, Ertelt JM, Jiang TT, Kinder JM, Xin L, Owens KJ, Jones HN, Way SS. CXCR3 blockade protects against *Listeria monocytogenes* infection-induced foetal wastage. *J Clin Invest.* (2015) 125(4):1713-25.
9. Masson E, Vidal C, Deschamps M, Bongain S, Thevenin C, Dupont I, Rietmulher D, Pouthier F, Mongaillard G, Chabod J, Ferrand C, Tiberghien P, Rebibou JM,

- 285 Incidence and risk factors of anti-HLA immunization after pregnancy. Hum  
 286 Immunol. (2013) Aug;74(8):946-51.  
 287
- 288 10. Meuleman T, van Beelen E, Kaaja RJ, van Lith JM, Claas FH, Bloemenkamp  
 289 KW, HLA-C antibodies in women with recurrent miscarriage suggests that antibody  
 290 mediated rejection is one of the mechanisms leading to recurrent miscarriage. J  
 291 Reprod Immunol. (2016) 116:28-34.
- 292 11. van Kampen CA, Versteeg-van der Voort Maarschalk MF, Langerak-Langerak J,  
 293 van Beelen E, Roelen DL, Claas FH, Pregnancy can induce long-persisting primed  
 294 CTLs specific for inherited paternal HLA antigens. Hum  
 295 Immunol. (2001) Mar;62(3):201-7
- 296 12. Lissauer, D., K. Piper, O. Goodyear, M. D. Kilby, and P. A. H. Moss, Fetal-  
 297 specific CD8+ cytotoxic T cell responses develop during normal human pregnancy  
 298 and exhibit broad functional capacity. J. Immunol. (2012) 189: 1072–80.
- 299 13. Piper, K. P., A. McLarnon, J. Arrazi, C. Horlock, J. Ainsworth, M. D. Kilby, W. L.  
 300 Martin, and P. Moss, Functional HY-specific CD8+ T cells are found in a high  
 301 proportion of women following pregnancy with a male fetus. Biol. Reprod. (2007) 76:  
 302 96–101.
- 303 14. Verdijk, R. M., A. Kloosterman, J. Pool, M. van de Keur, A. M. I. H. Naipal, A. G.  
 304 S. van Halteren, A. Brand, T. Mutis, and E. Goulmy, Pregnancy induces minor  
 305 histocompatibility antigen-specific cytotoxic T cells: implications for stem cell  
 306 transplantation and immunotherapy. Blood (2004) 103: 1961–4.
- 307 15. Verdijk, R. M., A. Kloosterman, J. Pool, M. van de Keur, A. M. I. H. Naipal, A. G.  
 308 S. van Halteren, A. Brand, T. Mutis, and E. Goulmy, Pregnancy induces minor  
 309 histocompatibility antigen-specific cytotoxic T cells: implications for stem cell  
 310 transplantation and immunotherapy. Blood (2004) 103.
- 311 16. van Halteren, A. G. S., E. Jankowska-Gan, A. Joosten, E. Blokland, J. Pool, A.  
 312 Brand, W. J. Burlingham, and E. Goulmy. Naturally acquired tolerance and  
 313 sensitization to minor histocompatibility antigens in healthy family members. Blood  
 314 (2009) 114.

17. Tilburgs, T., D. Schonkeren, M. Eikmans, N. M. Nagtzaam, G. Datema, G. M. Swings, F. Prins, J. M. van Lith, B. J. van der Mast, D. L. Roelen, S. A. Scherjon, and F. H. Claas. Human decidual tissue contains differentiated CD8<sup>+</sup> effector-memory T cells with unique properties. *J. Immunol.* (2010) 185: 4470–7.
18. Lissauer, D., S. A. Eldershaw, C. F. Inman, A. Coomarasamy, P. A. H. Moss, and M. D. Kilby, Progesterone promotes maternal-fetal tolerance by reducing human maternal T-cell polyfunctionality and inducing a specific cytokine profile. *Eur. J. Immunol.* (2015) 45: 2858–72.
19. Raghupathy, R., E. Al Mutawa, M. Makhseed, F. Azizieh, and J. Szekeres-Bartho, Modulation of cytokine production by dydrogesterone in lymphocytes from women with recurrent miscarriage. (2005) *BJOG* 112: 1096–101.
20. Chard, T., P. H. Craig, M. Menabawey, and C. Lee, Alpha interferon in human pregnancy. *Br. J. Obstet. Gynaecol.* (1986) 93: 1145–9.
21. Khan, N. U., A. Gibson, and A. K. Foulis, The distribution of immunoreactive interferon-alpha in formalin-fixed paraffin-embedded normal human foetal and infant tissues. *Immunology* (1990) 71: 230–5.
22. Bannert, N., and R. Kurth, Retroelements and the human genome: new perspectives on an old relation. *Proc. Natl. Acad. Sci. U. S. A.* (2004) 14572–9.
23. Tristem, M, Identification and characterization of novel human endogenous retrovirus families by phylogenetic screening of the human genome mapping project database. (2000) *J. Virol.* 74: 3715–30.
24. Blond, J. L., F. Besème, L. Duret, O. Bouton, F. Bedin, H. Perron, B. Mandrand, and F. Mallet, Molecular characterization and placental expression of HERV-W, a new human endogenous retrovirus family. *J. Virol.* (1999) 73: 1175–85.
25. McCoy, J. M., S. Mi, X. Lee, X. Li, G. M. Veldman, H. Finnerty, L. Racie, E. LaVallie, X.-Y. Tang, P. Edouard, S. Howes, and J. C. Keith, Syncytin is a captive retroviral envelope protein involved in human placentalmorphogenesis. *Nature* (2000) 403: 785–789.
26. Kämmerer, U., A. Germeyer, S. Stengel, M. Kapp, and J. Denner, Human

- 344 endogenous retrovirus K (HERV-K) is expressed in villous and extravillous  
345 cytotrophoblast cells of the human placenta. *J. Reprod. Immunol.* (2011) 91: 1–8.
- 346 27. Holder, B. S., C. L. Tower, K. Forbes, M. J. Mulla, J. D. Aplin, and V. M.  
347 Abrahams, Immune cell activation by trophoblast-derived microvesicles is mediated  
348 by syncytin 1. *Immunology* (2012) 136: 184–191.
- 349 28. Crespo ÂC, van der Zwan A, Ramalho-Santos J, Strominger JL, Tilburgs T,  
350 Cytotoxic potential of decidual NK cells and CD8+ T cells awakened by infections. *J*  
351 *Reprod Immunol.* (2017) Feb;119:85-90.
- 352 29. van Egmond A, van der Keur C, Swings GM, Scherjon SA, Claas FH, The  
353 possible role of virus-specific CD8(+) memory T cells in decidual tissue. *J Reprod*  
354 *Immunol.* (2016) Feb;113:1-8.
- 355 30. Crespo ÂC, Strominger JL, Tilburgs T. Expression of KIR2DS1 by decidual  
356 natural killer cells increases their ability to control placental HCMV infection. *Proc*  
357 *Natl Acad Sci U S A.* (2017) 113(52):15072-15077.  
358
- 359 31. Rowe, J. H., J. M. Ertelt, L. Xin, and S. S. Way, Pregnancy imprints regulatory  
360 memory that sustains anergy to fetal antigen. *Nature* (2012) 490: 102–6.
- 361 32. Aluvihare, V. R., M. Kallikourdis, and A. G. Betz, Regulatory T cells mediate  
362 maternal tolerance to the fetus. *Nat. Immunol.* (2004) 5: 266–71.
- 363 33. Tilburgs, T., D. L. Roelen, B. J. van der Mast, G. M. de Groot-Swings, C.  
364 Kleijburg, S. A. Scherjon, and F. H. Claas. Evidence for a selective migration of  
365 fetus-specific CD4+CD25bright regulatory T cells from the peripheral blood to the  
366 decidua in human pregnancy. *J. Immunol.* (2008) 180: 5737–45.
- 367 34. Chen T, Darrasse-Jèze G, Bergot AS, Courau T, Churlaud G, Valdivia  
368 K, Strominger JL, Ruocco MG, Chaouat G and Klatzmann D. Self-specific  
369 memory regulatory T cells protect embryos at implantation in mice. *J*  
370 *Immunol.* (2013) 191(5):2273-81.
- 371 35. Samstein, R. M., A. Arvey, S. Z. Josefowicz, X. Peng, A. Reynolds, R.  
372 Sandstrom, S. Neph, P. Sabo, J. M. Kim, W. Liao, M. O. Li, C. Leslie, J. A.  
373 Stamatoyannopoulos, and A. Y. Rudensky, Foxp3 exploits a pre-existent enhancer  
374 landscape for regulatory T cell lineage specification. *Cell* (2012) 151: 153–66.



36. Liang, S. C., Y. E. Latchman, J. E. Buhlmann, M. F. Tomczak, B. H. Horwitz, G. J. Freeman, and A. H. Sharpe, Regulation of PD-1, PD-L1, and PD-L2 expression during normal and autoimmune responses. *Eur. J. Immunol.* (2003) 33: 2706–16.
37. Tanaka, K., M. J. Albin, X. Yuan, K. Yamaura, A. Habicht, T. Murayama, M. Grimm, A. M. Waaga, T. Ueno, R. F. Padera, H. Yagita, M. Azuma, T. Shin, B. R. Blazar, D. M. Rothstein, M. H. Sayegh, and N. Najafian, PDL1 is required for peripheral transplantation tolerance and protection from chronic allograft rejection. *J. Immunol.* (2007) 179: 5204–10.
38. Topalian, S. L., F. S. Hodi, J. R. Brahmer, S. N. Gettinger, D. C. Smith, D. F. McDermott, J. D. Powderly, R. D. Carvajal, J. A. Sosman, M. B. Atkins, P. D. Leming, D. R. Spigel, S. J. Antonia, L. Horn, C. G. Drake, D. M. Pardoll, L. Chen, W. H. Sharfman, R. A. Anders, J. M. Taube, T. L. McMiller, H. Xu, A. J. Korman, M. Jure-Kunkel, S. Agrawal, D. McDonald, G. D. Kolli, A. Gupta, J. M. Wigginton, and M. Sznol. 2012. Safety, activity, and immune correlates of anti-PD-1 antibody in cancer. *N. Engl. J. Med.* (2012) 366: 2443–54.
39. Guleria, I., A. Khosroshahi, M. J. Ansari, A. Habicht, M. Azuma, H. Yagita, R. J. Noelle, A. Coyle, A. L. Mellor, S. J. Khoury, and M. H. Sayegh, A critical role for the programmed death ligand 1 in fetomaternal tolerance. *J. Exp. Med.* (2005) 202.
40. Wang, S., X. Zhu, Y. Xu, D. Zhang, Y. Li, Y. Tao, H. Piao, D. Li, and M. Du. Programmed cell death-1 (PD-1) and T-cell immunoglobulin mucin-3 (Tim-3) regulate CD4<sup>+</sup> T cells to induce Type 2 helper T cell (Th2) bias at the maternal-fetal interface. *Hum. Reprod.* (2016) 31: 700–11.
41. Chabtini, L., B. Mfarrej, M. Mounayar, B. Zhu, I. Batal, P. J. Dakle, B. D. Smith, O. Boenisch, N. Najafian, H. Akiba, H. Yagita, and I. Guleria, TIM-3 regulates innate immune cells to induce fetomaternal tolerance. *J. Immunol.* (2013) 190: 88–96.
42. Veras, E., R. J. Kurman, T.-L. Wang, and I.-M. Shih, PD-L1 Expression in Human Placentas and Gestational Trophoblastic Diseases. *Int. J. Gynecol. Pathol.* (2017). Mar;36(2):146-153
43. Wang, S.-C., Y.-H. Li, H.-L. Piao, X.-W. Hong, D. Zhang, Y.-Y. Xu, Y. Tao, Y. Wang, M.-M. Yuan, D.-J. Li, and M.-R. Du, PD-1 and Tim-3 pathways are associated

with regulatory CD8+ T-cell function in decidua and maintenance of normal pregnancy. (2015) *Cell Death Dis.* 6: e1738.

44. Old, LJ. Cancer is a somatic cell pregnancy. *Cancer Immun.* (2007) Nov 6;7:19.

45. Bianchi DW. Progress in the genetic analysis of fetal cells circulating in maternal blood. *Curr Opin Obstet Gynecol.* (1997) 9(2):121-5

46. Gammill HS, Guthrie KA, Aydelotte TM, Adams Waldorf KM, Nelson JL, Effect of parity on fetal and maternal microchimerism: interaction of grafts within a host? *Blood.* (2010) 116(15):2706-12.

47. Leavey, K., S. J. Benton, D. Gynspan, J. C. Kingdom, S. A. Bainbridge, and B. J. Cox, Unsupervised Placental Gene Expression Profiling Identifies Clinically Relevant Subclasses of Human Preeclampsia. *Hypertens.* (2016) 68: 137–47.

48. Kapur, P., D. Rakheja, A. M. Gomez, J. Sheffield, P. Sanchez, and B. B. Rogers, Characterization of Inflammation in Syphilitic Villitis and in Villitis of Unknown Etiology. *Pediatr. Dev. Pathol.* (2004) 7: 453–458.

49. Redline, R. W., and P. Patterson, Villitis of unknown etiology is associated with major infiltration of fetal tissue by maternal inflammatory cells. *Am. J. Pathol.* (1993) 143: 473–9.

50. Redline, R. W, Villitis of unknown etiology: noninfectious chronic villitis in the placenta. *Hum. Pathol.* (2007) 38: 1439–1446.

51. Tamblyn, J. A., D. M. Lissauer, R. Powell, P. Cox, and M. D. Kilby. The immunological basis of villitis of unknown etiology – Review. *Placenta* (2013) 34: 846–855.


52. Faas MM, de Vos P. Uterine NK cells and macrophages in pregnancy. *Placenta.* (2017) Aug;56:44-52.

ACCEPTED MANUSCRIPT

**Figure 1. Representation of maternal effector T cells within decidua and their potential functions.** These are likely to include (1) recognition of fetal antigen or virally-infected EVT through peptides on HLA-C alleles, (2) potential access to maternal blood vessels to limit systemic fetal chimerism and (3) potential to breach syncytiotrophoblast and damage fetal blood vessels

	Effector T cells within decidua	Effector T cells within maternal peripheral blood
Percentage of total T cell repertoire	~60%	~30%
Degree of differentiation	<b>more</b> differentiated (~40% CD27-CD28-)	<b>less</b> differentiated (~20% CD27-CD28-)
Pattern of cytokine production following stimulation with PMA/Ionomycin mitogen	<b>CD4+:</b> IFN $\gamma$ , 25%; IL-4, 5%  <b>CD8+:</b> IFN $\gamma$ , 60%; IL-4, 1.2%	<b>CD4+:</b> IFN $\gamma$ , 17%; IL-4, 2%  <b>CD8+:</b> IFN $\gamma$ , 41%; IL-4, 0.7%
Expression of checkpoint proteins	<b>CD4+</b> 43% PD-1+ <b>CD8+</b> 68% PD-1+	<b>CD4+</b> 20% PD-1+ <b>CD8+</b> 25% PD-1+
Frequency of fetal-specific T cells	Use of HLA-peptide multimers reveals <b>increased</b> numbers compared to blood	Rare – but potential role in <b>controlling fetal chimerism</b>
Differentially expressed genes	Increase in genes which mediate <b>interferon signaling response</b>	

Table 1. Comparison of the features of CD45RO+ effector T cells within decidua and maternal peripheral blood at term. Data from Powell et al, (Submitted).



**Figure 1. Representation of maternal effector T cells within decidua and their potential function.** These are likely to include (1) recognition of fetal antigen *or* virally infected EVT through peptides on HLA-C alleles, (2) potential access to maternal blood vessels to limit systemic fetal chimerism and (3) potential to breach damaged syncytiotrophoblast and damage fetal blood vessels

**Highlights**

**Maternal effector T cells within decidua: the adaptive immune response to pregnancy?**

**Lissauer D<sup>1</sup>, Kilby, MD<sup>1</sup>. and Moss, P<sup>2,3</sup>**

**Highlights**

**CD45RO<sup>+</sup> effector T cells comprise the majority of CD4<sup>+</sup> and CD8<sup>+</sup> T cells in decidua and are more highly differentiated than T cells in blood**

**Human effector T cells express CXCR3 which may guide cells to decidua**

**These cells include populations that can produce IFN $\gamma$  or IL-4**

**Microarray shows that decidual T cells demonstrate a transcriptional response to interferon signalling**

**T cells proliferate to cord blood indicating a response to fetal antigen and this is increased when T regulatory cells are removed.**